


Nested Gulf of Mexico Modeling with HYCOM



Patrick J. Hogan
Alan J. Wallcraft



Naval Research Laboratory
Stennis Space Center, MS

HYCOM Meeting
December 6-8, 2005
University of Miami, Miami, FL

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE DEC 2005		2. REPORT TYPE		3. DATES COVERED 00-00-2005 to 00-00-2005	
4. TITLE AND SUBTITLE Nested Gulf of Mexico Modeling with HYCOM			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,Stennis Space Center,MS,39529			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 25	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			


1/25° (~4 km) non-assimilative Nested Gulf of Mexico

- Initialized from January 1, 2000 1/12° Atlantic HYCOM
- Lateral boundary conditions from 1/12° Atlantic HYCOM
- Surface forcing is from 6-hourly/3-hourly NOGAPS (2000/2001)
- 20 layers in the vertical (bottom 5 from Atlantic discarded)
- 16 Rivers included as salinity flux
- GISS Mixed Layer submodel
- Bottom topography is from NRL-DBDB2
- Integrated over 2000-2001

Possible cross-shelf transport mechanisms

- Shelf break instabilities
- Shelf break flow reversals
- Eddy-eddy interactions
- Eddy-topography influence

1/25° Gulf of Mexico Model (~4 km)




Method of Characteristics used
To update the barotropic mode

20 gridpoint buffer zone for baroclinic
mode with e-folding time 1 to 10 days


Atlantic boundary data provided daily

1/25° (~4 km) Nested Gulf of Mexico



Snapshot of SSH
and SST on June, 13
2000

Lots of cyclonic
cold core eddies
(impact of 2x res.)




Local upwelling

1/25° Nested Gulf of Mexico HYCOM

Monthly Mean Sea Surface Height Year 2000

Evolution of Loop Current Eddy Shedding




- Loop Current Eddy sheds in August 2000
- Role of cyclones in Loop Current Eddy shedding evident

1/25° Nested Gulf of Mexico HYCOM

Monthly Mean Sea Surface Height Year 2001

Evolution of Loop Current Eddy Shedding



- Loop Current Eddy sheds 10 months later (July 2001)
- Detached eddy reattaches in several cases


1/25° Nested Gulf of Mexico HYCOM

red=east blue=west

July 27, 2000


red=north blue=south

Barotropic u-vel (cm/s) - 2000_208 (archive)



Barotropic u-velocity

Barotropic v-vel (cm/s) - 2000_208 (archive)




Barotropic v-velocity

**Initial eastward along-shelf break current
in geostrophic balance**


1/25° Nested Gulf of Mexico HYCOM

red=east blue=west August 12, 2000 (+ ~2 weeks) red=north blue=south

Barotropic u-vel (cm/s) - 2000_224 (archive)



Barotropic v-vel (cm/s) - 2000_224 (archive)




A reversal in the barotropic currents triggers a transition of the along-shelf break currents to flow onto the shelf

1/25° Nested Gulf of Mexico HYCOM


August 18, 2004 (+ ~1 week)

Barotropic u-vel (cm/s) - 2000_230 (archive)



red=east blue=west


Barotropic v-vel (cm/s) - 2000_230 (archive)



red=north blue=south

Significant cross-shelf flow exists after the reversal

1/25° Nested HYCOM forced by NOGAPS



Jet flowing eastward
along isobaths

The depth changes in
isopycnals across the current
serve as a potential energy
source for baroclinic eddies

Shelf break eddies associated
with baroclinic instability of the
along shelf-break current


April 10, 2000

red=east blue=west

1/25° Nested Gulf of Mexico HYCOM

Barotropic u-velocity


red=east blue=west



Meandering of the along-shelf jet

Barotropic v-velocity


red=north blue=south



Clockwise along-shelf break flow common in eastern Gulf

1/25° Nested Gulf of Mexico HYCOM



Jan 26 2001 SSH and Surface Currents



cyclones

- Cyclone cores are ~700m deep
- Robust shelf-break current associated with strong salinity gradient
- Loop Current penetration to ~28°N

Cross-section along 88.4°W




Red=east
Blue=west

1/25° Nested Gulf of Mexico HYCOM

Red=north
Blue=south



Feb 12, 2001 SSH and Surface Currents



Loop Current has migrated to NE
and is impinging on shelfbreak

Cyclone also impinging on shelfbreak

Cross-section along 27.9°N






- Doming of isohalines associated with cyclone
- Sharp shelfbreak front
- Intense northward subsurface jet

1/25° Nested Gulf of Mexico HYCOM

Red=north
Blue=south

March 01, 2001 SSH and Surface Currents






- Cyclone orbiting Loop Current Eddy,
- Loop Current Eddy breaching shelf break
- Southward flow enhanced by vortex compression?

Southward subsurface velocity maximum

1/25° Nested Gulf of Mexico HYCOM

Red=north
Blue=south

June 02, 2001 SSH and Surface Currents











3 months later cyclone hasn't migrated very far but is being steered by the shelf break

- Strange symmetry of LCE especially on western side
- Subsurface salinity max beneath LCE

January 05, 2002

Upwelling in the northern Gulf of Mexico




Layer 9

Northward velocity along shelf

Upwelling of cold water

Sensitivity of boundary forcing updating

Allows for long-term integrations over any timeframe



Monthly climatology formed from 1-day archives

Conclusions

Value of 2x resolution clear (cyclones, filaments, etc)

3 primary mechanisms of cross-shelf exchange:

- Along-shelf flow instabilities
- Along-shelf flow reversals
- Eddy-eddy interactions
- Influence of topography

Upwelling well represented (Yucatan, northern Gulf)

Climatological boundary forcing doesn't appear to
Degrade the solution

SUPPLEMENTAL SLIDES FOLLOW







Yucatan Channel Normal Velocity

0.08° ATL HYCOM 1-Year Mean

Observed Mean 8/1999-6/2000
(Abascal, et. al, 2001)



Note: section and sample period not identical



1/25° Free-Running Gulf of Mexico HYCOM




RMS SSH Variability

KPP




MY-2.5

GISS

2000



2001




KPP variability low in 2001


MY-2.5 variability low in 2000 and 2001

Need longer time series for meaningful statistics


layer=07 velocity mean: 2001.51-2001.59 [01.6H]



layer=20 velocity mean: 2001.51-2001.59 [01.6H]



layer=07 ke/mass mean: 2001.51-2001.59 [01.6H]



layer=20 ke/mass mean: 2001.51-2001.59 [01.6H]

